



TAROLA PLASTIC SURGERY  
NICHOLAS A. TAROLA, MD

---

## Blepharoplasty

**If unable to fully close lids, place eye ointment on inside of lower lid** or directly on eye before going to sleep or napping, beginning the day of surgery through the following 1-2 weeks, until able to fully close lids or otherwise instructed. Do not pull down eyelid when inserting ointment if you've had a lower blepharoplasty.

**Gently rinse eyes with eye saline solution** (Artificial Tears) upon waking through the first 1-2 weeks, until otherwise instructed. While awake, moisturize eyes with saline or eye drops as needed.

**Apply ice water soaked gauze** or clean washcloth to eyes for 20 minutes every two hours for the first 24 hours following surgery, and thereafter as needed for mild swelling.

**Avoid straining** or any activity that causes a feeling of pressure in the face and eyes for two weeks. No bending. Avoid sun exposure as much as possible, wear a hat and sunglasses when in direct sunlight.

**Do not take** any products containing **ibuprofen** (including Advil, Midol, etc.) in place of pain medicine for the first 24 hours after surgery.

**Some bruising and swelling is normal** as well as a small amount of oozing along the incision lines.

Please notify us if you are experiencing extreme swelling or bleeding from the incisions.

**Resuming social activities** and returning to work depend on the level of physical activity and public contact you require, as well as the swelling and bruising you may develop. The average patient goes back to work or social activities in 7 to 14 days.

### CALL THE OFFICE IMMEDIATELY IF YOU EXPERIENCE ANY OF THE FOLLOWING:

- **A high fever (over 101°), severe nausea and vomiting, continued dizziness or incoherent behavior, such as hallucinations**
- **Any pain that cannot be controlled by your pain medication**
- **Excessive bleeding or fluid seeping through the incisions**
- **Change in vision. If sudden loss of vision, go to nearest emergency room immediately**